

Not for Sale, Sale Into, Distribution, and/or Use in Nassau, Suffolk, Kings, and Queens Counties of New York State unless permitted under FIFRA Section 24(c) Special Local Need Registration.

CHLORANTRANILIPROLE	GROUP	28	INSECTICIDE
THIAMETHOXAM	GROUP	4A	INSECTICIDE



Insecticide

For control of listed insect pests of turfgrass and ornamental plants on golf courses, residential lawns, commercial grounds (i.e., office and shopping complexes, airports), parks, playgrounds, and athletic fields

Active Ingredients:	
Chlorantraniliprole*:	8.8%
Thiamethoxam**:	17.5%
Other Ingredients:	73.7%
Total:	100.0%

*CAS No. 500008-45-7

**CAS No. 153719-23-4

Acelepryn® Xtra is a suspension concentrate containing 1.67 lb of thiamethoxam and 0.835 lb chlorantraniliprole per gallon.

KEEP OUT OF REACH OF CHILDREN. CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional Precautionary Statements and Directions for Use inside booklet.

EPA Reg. No. 100-1680

EPA Est. 100-NE-001

SCP 1680A-L1 0623

2.5 gal
Net Contents



TABLE OF CONTENTS

1.0 FIRST AID

PRECAUTIONARY STATEMENTS

2.0 PRECAUTIONARY STATEMENTS

- 2.1 Hazards to Humans and Domestic Animals
- 2.2 Personal Protective Equipment (PPE)
- 2.3 User Safety Requirements
- 2.4 User Safety Recommendations
- 2.5 Environmental Hazards
 - 2.5.1 Groundwater Advisory
 - 2.5.2 Surface Water Advisory
 - 2.5.3 Pollinator Precautions
 - 2.5.4 Spray Drift Advisory
- 2.6 Physical or Chemical Hazards

DIRECTIONS FOR USE

3.0 USE RESTRICTIONS

4.0 PRODUCT INFORMATION

- 4.1 Foliar and Systemic Activity
- 4.2 Use Sites
- 4.3 Plant Tolerance
- 4.4 Integrated Pest Management (IPM)
- 4.5 Resistance Management Recommendations

5.0 APPLICATION DIRECTIONS

- 5.1 Methods of Application
- 5.2 Application Equipment
 - 5.2.1 Cleaning of Application Equipment
 - 5.2.2 Calibration of Application Equipment
- 5.3 Application Volume and Spray Coverage
- 5.4 Mixing Directions
 - 5.4.1 Acelepryn Xtra Alone
 - 5.4.2 Tank-Mix Precautions
 - 5.4.3 Tank-Mix Compatibility
 - 5.4.4 Acelepryn Xtra in Tank Mixtures
 - 5.4.5 Spray Additives
 - 5.4.6 Tank Mix Recommendations

6.0 APPLICATIONS TO TURFGRASS AND ORNAMENTAL PLANTS IN OUTDOOR LANDSCAPES

- 6.1 Turfgrass
- 6.2 Ornamental Plants in Outdoor Landscapes
 - 6.2.1 Foliar Applications
 - 6.2.2 Soil Applications for White Grubs
 - 6.2.3 Soil Treatments for Above-Ground Pests
- 6.3 Individual Fire Ant Mound Treatment

7.0 STORAGE AND DISPOSAL

8.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

1.0 FIRST AID

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
NOTE TO PHYSICIAN There is no specific antidote if Acelepryn Xtra is ingested. Induce emesis or lavage stomach. Treat symptomatically.	
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
HOTLINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

2.0 PRECAUTIONARY STATEMENTS

2.1 Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

2.2 Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate; butyl rubber ≥ 14 mils; nitrile rubber ≥ 14 mils; neoprene rubber ≥ 14 mils; polyvinyl chloride (PVC) ≥ 14 mils; or Viton™ ≥ 14 mils
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

2.3 User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

2.4 User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

2.5 Environmental Hazards

This pesticide is toxic to wildlife and highly toxic to aquatic invertebrates, oysters, and shrimp.

This product is highly toxic to bees exposed to direct treatment on blooming plants or weeds and may cause possible effects to pollinators from exposure to translocated residues in blooming plants. Do not apply this product or allow it to drift to blooming plants or weeds while bees are foraging in or adjacent to the treatment area.

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or disposing of equipment wash-water or rinsate.

2.5.1 GROUNDWATER ADVISORY

This product has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into the groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

2.5.2 SURFACE WATER ADVISORY

This product may impact surface water quality due to spray drift and runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several months after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of thiamethoxam from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours. (See manual at the following Internet address:

<https://www.nrcs.usda.gov/conservation-basics/natural-resource-concerns/water/>)

2.5.3 POLLINATOR PRECAUTIONS

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications or contact with residues on plant surfaces after foliar applications
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <https://pesticidestewardship.org/pollinator-protection/>

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: <https://aapco.org/2015/07/28/resources-2/>. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

2.5.4 SPRAY DRIFT ADVISORY

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS (i.e., 15 mph or greater). Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

2.6 Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read and understand the entire label directions, restrictions, and precautions before using this product. All applications of Acelepryn Xtra must be made only according to the use directions on this label or in EPA approved labeling (Supplemental Labels, Special Local Need Registrations, FIFRA Section 18 exemptions).

FOR NON-AGRICULTURAL USES ONLY



- Do not apply Acelepryn Xtra while bees are foraging. Do not apply Acelepryn Xtra to plants that are flowering. Only apply after all flower petals have fallen off.

For any requirements specific to your state or tribe, consult the state or tribal agency responsible for pesticide regulation.

Notify state and/or federal authorities and Syngenta immediately if you observe any adverse effects due to use of this product.

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY AND/OR POOR INSECT CONTROL.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides, 40 CFR Part 170. The WPS applies when this product is used to produce agricultural plants on ornamental nurseries or greenhouses, sod farms, or forests.

Professional applications to golf courses, residential, industrial, and commercial lawns, and sports fields are not within the scope of the Worker Protection Standard.

Do not enter or allow others to enter the treated area until sprays have dried.

3.0 USE RESTRICTIONS

- DO NOT apply more than 20 fluid ounces of Acelepryn Xtra per acre per calendar year or:
 - greater than 0.50 lb of chlorantraniliprole-containing products
 - greater than 0.265 lb of thiamethoxam-containing products
- DO NOT let people and pets enter the treated area until treatment has dried.
- DO NOT allow this product to contact plants in bloom while bees are foraging in the treatment area.
- DO NOT apply this product, by any application method, to linden, basswood, or *Tilia* species.
- DO NOT use on plants being grown for sale, seed production, or research purposes.
- DO NOT use in the commercial nursery production of ornamental plants.
- DO NOT use in greenhouses.
- DO NOT apply this product through any type of irrigation system.
- DO NOT apply this product with aerial application equipment.
- DO NOT apply by ground equipment within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.

- Not for Sale, Sale Into, Distribution and/or Use in Nassau, Suffolk, Kings, and Queens Counties of New York State unless permitted under FIFRA Section 24(c) Special Local Need Registration.
- DO NOT apply Acelepryn Xtra to plants that are flowering. Only apply after all flower petals have fallen off.

The following restrictions are required to permit use of Acelepryn Xtra in the State of New York:

- This product is classified as restricted in New York State.
- Golf course greens and tee boxes: DO NOT apply this product within 25 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).
- Golf course fairways: DO NOT apply this product within 50 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).
- For all other application sites, DO NOT apply this product within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).
- In New York State, do not exceed a total of 0.188 lb ai of thiamethoxam-containing products per acre per growing season (for outdoor applications).

4.0 PRODUCT INFORMATION

4.1 Foliar and Systemic Activity

Acelepryn Xtra contains systemic insecticides that are translocated upward into the plant from root uptake when applied to the soil. To be effective, apply Acelepryn Xtra where the root system of the target plant can readily absorb the active ingredients.

4.2 Use Sites

For use on ornamental plants and turfgrasses on golf courses, residential lawns, commercial grounds (i.e., office and shopping complexes, airports), parks, playgrounds, and athletic fields. Not for use on plants or turfgrasses being grown for sale or other commercial use; commercial seed production; or for research purposes. Not for use in greenhouses.

4.3 Plant Tolerance

Certain plant species or cultivars may be sensitive to the final spray solution. If local experience is not available, test Acelepryn Xtra alone or the mixture to be applied (including adjuvants) to a small number of plants and observe for phytotoxicity for at least one week before treating the entire planting to ensure plant safety.

4.4 Integrated Pest Management (IPM)

Some insect pests are known to develop resistance to products after repeated use. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies established for the crop and use area. Syngenta encourages responsible product stewardship to ensure effective long-term control of the insects on the label.

4.5 Resistance Management Recommendations

CHLORANTRANILIPROLE	GROUP	28	INSECTICIDE
THIAMETHOXAM	GROUP	4A	INSECTICIDE

For resistance management, Acelepryn Xtra contains Group 28 and Group 4A insecticides. Any insect population may contain individuals naturally resistant to Acelepryn Xtra and other Group 28 and 4A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of Acelepryn Xtra or other Group 28 and 4A insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classifications are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.

- Adopt an integrated pest management program for insecticide use that includes pest monitoring, uses historical information related to pesticide use, record keeping, and which considers cultural, biological, and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance, contact Syngenta at 1-866-SYNGENT(A) (866-796-4368).

Syngenta encourages responsible product stewardship to ensure effective long-term control of the insects on this label.

5.0 APPLICATION DIRECTIONS

5.1 Methods of Application

Apply Acelepryn Xtra to turf and ornamental plant foliage or to the soil by ground equipment only as specified in **Sections 6.1 to 6.3**.

Make foliar applications with the following ground application equipment: boom sprayers and low pressure, handheld-application equipment (hand wand, backpack and mechanical hand gun sprayers).

Make soil and mound applications by soil injection, soil drenches, and broadcast, banded and directed sprays using boom and low pressure, handheld-application equipment (hand wand, backpack and mechanical hand gun sprayers).

5.2 Application Equipment

5.2.1 CLEANING OF APPLICATION EQUIPMENT

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all application equipment to reduce the risk of forming hardened deposits that can become difficult to remove. Drain application equipment. Thoroughly rinse application equipment and flush hoses, boom, and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean equipment near wells, water sources, or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

5.2.2 CALIBRATION OF APPLICATION EQUIPMENT

Proper maintenance and calibration of spraying equipment are essential for optimal insect pest control. If you have questions about calibration, contact a State Extension Service specialist, the equipment manufacturer, or other experts.

5.3 Application Volume and Spray Coverage

Acelepryn Xtra must be diluted with water before application. Apply in a volume of water that provides good coverage of the foliage or soil, but does not result in run-off or leaching, as specified in the Directions for Use in **Section 6.0**.

5.4 Mixing Directions

5.4.1 ACELEPRYN XTRA ALONE

Prepare only the amount of spray mixture that is needed for the immediate application. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area. Keep product container tightly closed when not in use.

1. Fill sprayer tank with 1/2 of the required amount of water
2. Add the required amount of Acelepryn Xtra directly to the sprayer tank with the mechanical or hydraulic agitator running. Do not use air agitation.
3. Continue agitation while adding the remainder of the water.
4. Begin application of the solution after Acelepryn Xtra has completely dispersed into the mix water.
5. Maintain agitation until all of the mixture has been applied.

5.4.2 TANK-MIX PRECAUTIONS

Acelepryn Xtra may be tank-mixed with other pesticides. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Do not mix this product with any product which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

5.4.3 TANK-MIX COMPATIBILITY

The physical compatibility of Acelepryn Xtra will vary with different sources of pesticide products and local cultural practices. To ensure the physical compatibility of the mixture, prepare a mix on a small scale (such as a pint or quart jar) using the proper proportions of pesticides and water. Add different formulation-type tank mix partners in the sequence indicated below in 5.4.4. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible.

5.4.4 ACELEPRYN XTRA IN TANK MIXTURES

Add 1/2 of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. Add different formulation-type tank mix partners in the sequence indicated below. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all the mixture has been applied.

1. Water-soluble bags
2. Water-dispersible granules
3. Wettable powders
4. Acelepryn Xtra (this product) and other water-based suspension concentrates
5. Water-based soluble concentrates
6. Emulsifiable concentrates
7. Adjuvants, surfactants, oils
8. Soluble fertilizers
9. Drift retardants

5.4.5 SPRAY ADDITIVES

When an adjuvant is to be used with this product, the use of an adjuvant that meets the standards of the Council of Producers and Distributors of Agrotechnology (CPDA) adjuvant certification is recommended.

When making foliar applications to hard-to-wet foliage, such as holly, pine, or ivy, the addition of a spreader/sticker is recommended.

5.4.6 TANK MIX RECOMMENDATIONS

Acelepryn Xtra may be tank mixed with a nematocide to improve nematode control and enhance turf quality.

When Acelepryn Xtra is used to control parasitic nematodes on golf course turf, a fungicide treatment is recommended in combination with Acelepryn Xtra plus a nematocide to reduce fungal infections following nematode feeding.

6.0 APPLICATIONS TO TURFGRASS AND ORNAMENTAL PLANTS IN OUTDOOR LANDSCAPES

- Apply Acelepryn Xtra as a broadcast, foliar, soil, or mound treatment to control listed insect pests of turf and ornamental plants grown on golf courses, residential lawns, commercial grounds (i.e., office and shopping complexes, airports), parks, playgrounds, and athletic fields. See **Tables 6.1 to 6.3** for specific Directions for Use.
- For maximum residual control or heavy pest pressure, apply at highest listed rate.
- Apply as a broadcast spray using a boom sprayer, low pressure handheld application equipment (hand-wand, backpack, and mechanical hand-gun sprayers).
- Apply in a volume of water sufficient to uniformly cover the area being treated (a minimum of 1.5 - 5 gallons per 1,000 square feet).
- Use properly calibrated application equipment that will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off-target drift.

6.1 Turfgrass

Use Sites: Turfgrasses on golf courses, residential lawns, commercial grounds (i.e., office and shopping complexes, airports), parks, playgrounds, and athletic fields			
Target Pests	Product Use Rate	Application Timing	Use Directions
Turf Caterpillars (including armyworms, cutworms, and sod webworms)	4 - 20 fl oz per acre (0.052 - 0.26 lb thiamethoxam and 0.026 - 0.13 lb chlorantraniliprole) or 1.0 - 4.6 fl oz per 10,000 sq ft (0.013 - 0.06 lb thiamethoxam and 0.0065 - 0.03 lb chlorantraniliprole)	Make application when populations are first observed.	If the area being treated is maintained at a mowing height of greater than one inch, then use the higher rates within the rate range to control heavy pressure. To ensure optimum control, for turf caterpillars, delay watering (irrigation) or mowing for 24 hours after application.

continued...

6.1 Turfgrass (continued)

Target Pests	Product Use Rate	Application Timing	Use Directions
White Grubs (Larvae of): (including <i>Aphodius</i> spp., Asiatic garden beetle, black turfgrass <i>Ataenius</i> *, European chafer, green June beetle, Japanese beetle, May/June beetles (<i>Phyllophaga</i> spp.), northern masked chafer, oriental beetle, southern masked chafer, sugarcane grub)	10 – 20 fl oz per acre (0.13 – 0.26 lb thiamethoxam and 0.065 – 0.13 lb chlorantraniliprole) or 2.3 – 4.6 fl oz per 10,000 sq ft (0.03 – 0.06 lb thiamethoxam and 0.015 – 0.03 lb chlorantraniliprole)	Apply Acelepryn Xtra from early April to early September for preventative and early remedial control of the listed white grub species infesting turfgrass. The need for an application may be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. *Applications targeting black turfgrass <i>Ataenius</i> larvae should be applied before the eggs of the first-generation hatch. In most areas, applications should be made during May to ensure control of the first-generation larvae. This application will control both first and second generation black turfgrass <i>ataenius</i> larvae.	Irrigate within 24 hours of application if sufficient rainfall (1/4-1/2 inch of water) does not occur. The high rate for white grub control may be required for early season (April and May) applications where a long residual is required or in late August or early September when less sensitive mid-instar grubs are present at the time of application.
Annual bluegrass weevil	10 – 20 fl oz per acre (0.13 – 0.26 lb thiamethoxam and 0.065 – 0.13 lb chlorantraniliprole) or 2.3 – 4.6 fl oz per 10,000 sq ft (0.03 – 0.06 lb thiamethoxam and 0.015 – 0.03 lb chlorantraniliprole)	Apply Acelepryn Xtra when overwintered adult annual bluegrass weevils are observed in late April or early May to prevent damage from first generation larvae in late-May and June.	Use the higher rates within the rate range after mid-May to control heavy Annual Bluegrass Weevil pressure
Billbugs	10 – 20 fl oz per acre (0.13 – 0.26 lb thiamethoxam and 0.065 – 0.13 lb chlorantraniliprole) or 2.3 – 4.6 fl oz per 10,000 sq ft (0.03 – 0.06 lb thiamethoxam and 0.015 – 0.03 lb chlorantraniliprole)	Apply Acelepryn Xtra when overwintered adult billbugs are first observed. This will usually occur in late April or early May in regions with cool-season turfgrasses.	
European Crane Fly	10 – 20 fl oz per acre (0.13 – 0.26 lb thiamethoxam and 0.065 – 0.13 lb chlorantraniliprole) or 2.3 – 4.6 fl oz per 10,000 sq ft (0.03 – 0.06 lb thiamethoxam and 0.015 – 0.03 lb chlorantraniliprole)	Apply Acelepryn Xtra (prior to egg hatch) between September and November to control European crane fly larvae in turfgrass. The high rate may be required to achieve control when applications are made in November.	

Target Pests	Product Use Rate	Application Timing	Use Directions
Chinch bugs	10 – 20 fl oz per acre (0.13 – 0.26 lb thiamethoxam and 0.065 – 0.13 lb chlorantraniliprole) or 2.3 – 4.6 fl oz per 10,000 sq ft (0.03 – 0.06 lb thiamethoxam and 0.015 – 0.03 lb chlorantraniliprole)	For optimum control apply when young nymphs are first observed.	
Flea beetles Greenbugs Leafhoppers Spittlebugs	15 – 20 fl oz per acre (0.2 – 0.26 lb thiamethoxam and 0.1 – 0.13 lb chlorantraniliprole) or 3.4 – 4.6 fl oz per 10,000 sq ft (0.044 – 0.06 lb thiamethoxam and 0.022 – 0.03 lb chlorantraniliprole)	For optimum control, make application when populations are first observed. Acelepryn Xtra will provide control of two-lined spittlebug when applications are made in spring or summer.	
Ants (excluding Carpenter, Harvester, and Pharaoh ants)	15 – 20 fl oz per acre (0.2 – 0.26 lb thiamethoxam and 0.1 – 0.13 lb chlorantraniliprole) or 3.4 – 4.6 fl oz per 10,000 sq ft (0.044 – 0.06 lb thiamethoxam and 0.022 – 0.03 lb chlorantraniliprole)	To control ants, treat when ant mounds are first observed.	To achieve optimum control on green and tee surfaces, treat the affected area plus a surrounding 30-foot buffer. For fairways and roughs, treat at least 1.5 times the infested area. Irrigate within 24 hours of application if sufficient rainfall (1/4-1/2 inch of water) does not occur. For additional activity, utilize with another insecticide such as lambda-cyhalothrin in an insect control spray program. Consult the other insecticide label for use instructions and sites applicability.
Mole crickets (suppression*)	15 – 20 fl oz per acre (0.2 – 0.26 lb thiamethoxam and 0.1 – 0.13 lb chlorantraniliprole) or 3.4 – 4.6 fl oz per 10,000 sq ft (0.044 – 0.06 lb thiamethoxam and 0.022 – 0.03 lb chlorantraniliprole)	To suppress damage, treat from first egg hatch to peak egg hatch (mid-summer to control 3rd and 4th instar larvae). Acelepryn Xtra provides suppression of mole crickets on turfgrass.	

continued...

6.1 Turfgrass (continued)

APPLICATION METHOD AND INFORMATION
<ul style="list-style-type: none"> Apply Acelepryn Xtra as a broadcast application to the listed turfgrass sites. Use a minimum water volume of 15-50 gallons/10,000 square feet to uniformly distribute this product over the area being treated and to adequately move the active ingredients into the thatch layers. For turf with heavy thatch (more than 0.75 inches) use the higher rates within the specified rate range. Make applications to turfgrass sites as broadcast sprays using a boom sprayer, low pressure handheld application equipment (hand-wands, backpack, and mechanical handgun sprayers). *Note: Suppression can mean either erratic control, ranging from good to poor, or a consistent level of control below that generally acceptable for commercial purposes.
USE RESTRICTIONS
<ol style="list-style-type: none"> Maximum Single Application Rate: <ol style="list-style-type: none"> Do not apply more than 20 fl oz of Acelepryn Xtra per acre per application. Maximum Annual Application Rate: <ol style="list-style-type: none"> Do not apply more than 20 fl oz of Acelepryn Xtra per acre per calendar year, or: <ol style="list-style-type: none"> greater than 0.50 lb ai per acre per calendar year from any chlorantraniliprole-containing products greater than 0.265 lb ai per acre per calendar year from any thiamethoxam-containing products Do not apply more than once every 7 days. Do not enter or allow others to enter treated areas until sprays have dried. DO NOT apply this product through any type of irrigation system.

6.2 Ornamental Plants in Outdoor Landscapes

6.2.1 FOLIAR APPLICATIONS

Use Sites: Ornamentals in outdoor landscapes on golf courses, residential lawns, commercial grounds (i.e., office and shopping complexes, airports), parks, playgrounds, and athletic fields			
Evergreens, including conifers	Ornamental trees ¹		
Flowering plants ¹	Shrubs ¹		
Foliage plants (non-food)			
Ground covers ¹			
Target Pests	Product Use Rate	Application Timing	Use Directions
Ants, excluding harvester, carpenter, and Pharaoh ants Aphids	2.4 – 4.8 fl oz per 100 gallons (0.031-0.062 lb thiamethoxam and 0.016-0.031 lb chlorantraniliprole)	Apply preventatively or after pests have been observed. Repeat treatment to maintain control using the higher listed application rates as pest pressure and foliage area increases.	Mix Acelepryn Xtra with the required amount of water and apply as a full-coverage foliar spray. For hard-to-wet foliage such as holly, pine, or ivy, the addition of a spreader/sticker is recommended.
Adelgids, including Hemlock Woolly Adelgid Lace Bugs Leaf Beetles Leaf-feeding caterpillars, such as bagworms and tussock moth caterpillars, including whitemarked tussock moth Leafhoppers Leafminer Leaf gall-forming insects Japanese beetle adults Mealybugs Midges Plant Bug Psyllids, including Asian Citrus Psyllid Sawflies Soft Scales Thrips – foliar feeding Whiteflies	4 – 16 fl oz per 100 gallons (0.052-0.208 lb thiamethoxam and 0.026-0.104 lb chlorantraniliprole)	For gall makers, apply during egg laying of the generation being targeted. For leafhoppers, plant bug, and aphids, make the first application when populations are first observed.	

Target Pests	Product Use Rate	Application Timing	Use Directions
Black vine weevil (adult) ²	9.6 – 16 fl oz per 100 gallons (0.125-0.208 lb thiamethoxam and 0.062-0.104 lb chlorantraniliprole)	For black vine weevil, apply to foliage of plants when adult feeding damage is first observed. Make only one application per generation.	
APPLICATION METHOD AND INFORMATION			
<ul style="list-style-type: none"> Apply Acelepryn Xtra as a broadcast or directed application to the listed ornamental plants. Use a minimum water volume of 15-50 gallons per 10,000 square feet to uniformly distribute this product over the area being treated. Make applications by low-pressure handheld application equipment (hand-wands, backpack, and mechanical handgun sprayers). 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> Maximum Single Application Rate: <ol style="list-style-type: none"> Do not apply more than 20 fl oz of Acelepryn Xtra per acre per application. Maximum Annual Application Rate: <ol style="list-style-type: none"> Do not apply more than 20 fl oz of Acelepryn Xtra per acre per calendar year, or: <ol style="list-style-type: none"> greater than 0.5 lb ai per acre per calendar year from any chlorantraniliprole-containing products greater than 0.26 lb ai per acre per calendar year from any thiamethoxam-containing products Do not apply more than once every 7 days. Do not enter or allow others to enter the treated area until sprays have dried. Do not apply Acelepryn Xtra to plants that are flowering. Only apply after all flower petals have fallen off. ¹Refer to Protection of Pollinators in Section 2.5.3 ²Not for use in California 			



6.2.2 SOIL APPLICATIONS FOR WHITE GRUBS

Use Sites: Ornamentals in outdoor landscapes on golf courses, residential lawns, commercial grounds (i.e., office and shopping complexes, airports), parks, playgrounds, and athletic fields			
Evergreens, including conifers	Ornamental trees ¹		
Flowering plants ¹	Shrubs ¹		
Foliage plants			
Ground covers ¹			
Target Pests	Product Use Rate	Application Timing	Use Directions
White grubs, including Asiatic garden beetle, European chafer, green June beetle, Japanese beetle, May/June beetles (<i>Phyllophaga</i> spp.), northern masked chafer, oriental beetle, and southern masked chafer	16 – 20 fl oz per acre (0.208-0.260 lb thiamethoxam and 0.104-0.130 lb chlorantraniliprole) or 3.7 – 4.6 fl oz per 10,000 sq ft (0.048-0.060 lb thiamethoxam and 0.024-0.030 lb chlorantraniliprole)	Apply up to 45 days before the historical peak of adult flight to 2nd instar grub of the species being targeted. For optimum control, treat from peak flight to peak egg hatch. To control grubs, make the application from egg hatch to second instar (grubs less than half their full size.)	Apply in sufficient water to uniformly cover the area being treated. Irrigate within 24 hours of application if sufficient rainfall (1/4-1/2 inch of water) does not occur.
APPLICATION METHOD AND INFORMATION			
<ul style="list-style-type: none"> Make applications as a broadcast application through low-pressure handheld application equipment (hand-wands, backpack, and mechanical handgun sprayers). Use sufficient water volume (15 – 50 gallons/10,000 square feet) to uniformly distribute Acelepryn Xtra over the area being treated. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> Maximum Single Application Rate: <ol style="list-style-type: none"> Do not apply more than 20 fl oz of Acelepryn Xtra per acre per application. Maximum Annual Application Rate: <ol style="list-style-type: none"> Do not apply more than 20 fl oz of Acelepryn Xtra per acre per calendar year, or: <ol style="list-style-type: none"> greater than 0.50 lb ai per acre per calendar year from any chlorantraniliprole-containing products greater than 0.265 lb ai per acre per calendar year from any thiamethoxam-containing products Do not apply more than once every 7 days. Do not enter or allow others to enter the treated area until sprays have dried. ¹Refer to Protection of Pollinators in Section 2.5.3 			



6.2.3 SOIL TREATMENTS FOR ABOVE-GROUND PESTS

Use Sites: Ornamentals in outdoor landscapes on golf courses, residential lawns, commercial grounds (i.e., office and shopping complexes, airports), parks, playgrounds, and athletic fields			
Evergreens, including conifers Foliage plants	Ornamental trees ²	Palms Shrubs ²	
Target Pests	Product Use Rate ¹	Application Timing	Use Directions
Lace bugs Aphids, including apple aphid	<p>Trees: 0.125 - 0.17 fl oz per one inch of tree trunk diameter (DBH)</p> <p>Shrubs: 0.125 - 0.17 fl oz per foot of height of shrub height</p> <p>Hedge Row: 0.125 - 0.17 fl oz per foot of height of shrub height (< 6' tall) per plant</p>	<p>Apply 1 to 4 weeks before pest has historically been observed to achieve best control.</p> <p>When making soil applications to plants with woody stems, systemic activity will be delayed until the active ingredient is translocated throughout the plant. This delay could be 60 days or longer for larger plants.</p>	<p>Soil Injection Method for Trees: <i>Basal System:</i> Apply the proper rate among 4 evenly spaced injection sites located within a foot of each tree. <i>Grid System:</i> Apply the proper rate among injection sites on a grid pattern with 1.5-2.5 feet centers extending to the drip line of the tree.</p> <p>Soil Drench Method for Trees and Shrubs: Apply the proper rate uniformly around the base of the tree/shrub within 1-2 feet of the trunk/base. Use a minimum of 16 fluid ounces of solution per inch DBH or per foot of shrub height.</p> <p>For multiple plants (hedge row), determine the number of plants per hedge row, multiply by the average height and rate to be applied per foot of height.</p> <p>To be effective, apply Acelepryn Xtra at the proper rate uniformly within 1-3 feet of the root flare of trees and shrubs. Use a minimum of 16 fluid ounces solution to reach the root zone for optimal systemic uptake.</p>
Birch leafminer Rhododendron lace bug Hemlock woolly adelgid Magnolia scale False Oleander Scale	<p>Trees: 0.17 fl oz per one inch of tree trunk diameter (DBH)</p> <p>Shrubs: 0.17 fl oz per foot of height of shrub height</p> <p>Hedge Row: 0.17 fl oz per foot of height of shrub height (< 6' tall) per plant</p>		
APPLICATION METHOD AND INFORMATION			
<ul style="list-style-type: none"> ¹Amount of Acelepryn Xtra per inch of diameter at breast height (DBH) for trees or per foot of height for shrubs. (0.125 fl oz product = 0.0016 lb thiamethoxam and 0.0008 lb chlorantraniliprole, and 0.170 fl oz product = 0.0022 lb thiamethoxam and 0.0011 fl oz chlorantraniliprole) Make applications by soil injection or soil drench. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> Maximum Single Application Rate: <ol style="list-style-type: none"> Do not apply more than 20 fl oz of Acelepryn Xtra per acre per application. Maximum Annual Application Rate: <ol style="list-style-type: none"> Do not apply more than 20 fl oz of Acelepryn Xtra per acre per calendar year, or: <ol style="list-style-type: none"> greater than 0.50 lb ai per acre per calendar year from any chlorantraniliprole-containing products greater than 0.265 lb ai per acre per calendar year from any thiamethoxam-containing products Do not apply more than once every 7 days. Do not enter or allow others to enter the treated area until sprays have dried. ² Refer to Protection of Pollinators in Section 2.5.3 			



6.3 INDIVIDUAL FIRE ANT MOUND TREATMENT

Use Sites: Golf Courses, Lawns, and Recreational Turf Areas (parks, playgrounds, and athletic fields)			
Target Pests	Product Use Rate	Application Timing	Use Directions
Fire ants (<i>Solenopsis</i> spp.)	1.25 - 3.75 fl oz per 10 gallons (0.016-0.049 lb thiamethoxam and 0.008-0.024 lb chlorantraniliprole)	Make a single drench application when mounds appear to be active.	<p>Thoroughly mix solution and apply directly to mounds. For optimum control of small ant mounds (<6 inches in diameter at the surface), apply one gallon of the drench solution per mound.</p> <p>For optimum control of larger ant mounds, apply 2-3 gallons of the drench solution per mound.</p> <p>Direct the drench application at the center of the mound and include a 6-inch diameter circle around the center of the mound. Do not apply less than 0.5 gallon or more than 3 gallons of drench solution per mound.</p>
APPLICATION METHOD AND INFORMATION			
<ul style="list-style-type: none"> Make application by drench application using low pressure handheld application equipment (hand-wands, backpack, and mechanical handgun sprayers). 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> Maximum Single Application Rate: <ol style="list-style-type: none"> Do not apply more than 20.0 oz of Acelepryn Xtra per acre per application. Maximum Annual Application Rate: <ol style="list-style-type: none"> Do not apply more than 20.0 oz of Acelepryn Xtra per acre per calendar year, or: <ol style="list-style-type: none"> greater than 0.50 lb ai per acre per calendar year from any chlorantraniliprole-containing products greater than 0.265 lb ai per acre per calendar year from any thiamethoxam-containing products Do not apply more than once every 7 days. Do not enter or allow others to enter treated areas until sprays have dried. 			

7.0 STORAGE AND DISPOSAL

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Do not subject to temperatures below 32 degrees F. Store product in original container only in a location inaccessible to children and pets. Do not contaminate water, other pesticides, fertilizer, food, or feed in storage.

Pesticide Disposal

Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling HDPE Plastic (less than or equal to 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¹/₄ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

8.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and of Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

Acelepryn®, the ALLIANCE FRAME, the Syngenta Logo, and the PURPOSE ICON are Trademarks of a Syngenta Group Company

Viton™ is a registered trademark of the Chemours Company FC, LLC.

©2023 Syngenta

For non-emergency (e.g., current product information), call Syngenta Crop Protection at 1-866-796-4368.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1680A-L1 0623